

Resonance interaction between two entangled atoms in a photonic bandgap environment

Valentina Notararigo*

Department of Physics and Astronomy, University College London, Gower Street, London WC1E 6BT, UK

Roberto Passante[†] and Lucia Rizzuto[‡]

*Dipartimento di Fisica e Chimica, Università degli Studi di Palermo and CNISM, Via Archirafi 36, I-90123 Palermo, Italy
and INFN, Laboratori Nazionali del Sud, I-95123 Catania, Italy*

We consider the resonance interaction energy and force between two identical atoms, one in an excited state and the other in the ground state, interacting with the quantum electromagnetic field in the vacuum state and placed in a photonic-bandgap environment with a quadratic dispersion relation. Such an environment could be a coupled cavities array or a photonic crystal in the effective mass approximation, for example. The two atoms are prepared in their symmetric or antisymmetric entangled state; the atomic excitation is thus delocalized among the atoms. This problem is also strictly related to the coherent resonant energy transfer between two atoms. We consider the case of an atomic transition frequency inside the photonic gap and in proximity of its edge. We discuss both a one-dimensional and a three-dimensional case. In both cases, we find that the resonance interaction decays faster with the distance compared to the free-space case. In the one-dimensional case, it asymptotically decays as $1/r$, while in free space it has a pure oscillatory dependence; in the isotropic three-dimensional case, the resonance interaction asymptotically scales as $1/r^2$, compared with the $1/r$ dependence in free space. We argue that, although the resonance force can be reduced by the presence of the bandgap environment in the very long-distance regime (at intermediate distances it can be even increased), the configuration considered can be suitable for the observation of the resonance interaction because spontaneous emission is strongly suppressed by the environment and the correlated state is thus preserved by the decoherence effect due to its spontaneous decay.

PACS numbers: 12.20.Ds, 42.50.Ct

The resonance interaction energy and the energy transfer between two atoms coupled with the electromagnetic field in the vacuum state, one in the ground state and the other in the excited state, have recently received much attention in the literature in different research fields [1–4]. For example, coherent energy transfer has been investigated in quantum biology, where it is supposed to play an important role in the photosynthesis process [5, 6]. From a quantum-electrodynamical point of view, the resonance interaction is due to the exchange of a real or virtual photon between the two atoms.

If the two atoms are in a factorized state, their radiation-mediated interaction energy is a fourth-order process in the electric charge [7, 8], and controversial results exist in the literature concerning with the presence of spatially oscillating terms [9–12]. The interaction is a second-order effect when two identical atoms are prepared in a symmetric or antisymmetric entangled state [1, 8, 13]. The resonance interaction is particularly relevant because this interaction is much more intense than dispersion interactions (van der Waals/Casimir-Polder, both for ground-state and excited atoms) and Casimir-Polder interactions between a ground-state atom and a mirror, and comparable to the atom-surface interaction for

excited atoms [14]. The Casimir-Polder interaction between an excited atom and a surface has been recently measured directly [15, 16]. From a physical point of view, it is a second-order effect because the correlated state has nonvanishing dipole-dipole correlations, while in the case of dispersion interactions correlated dipole moments must be induced by vacuum fluctuations [17]. Moreover, the resonance interaction is of very long range, scaling in free space as r^{-1} , with space oscillations for distances larger than a characteristic atomic transition wavelength (three-dimensional case) [1], whilst in the same limit it is just an oscillating function in the one-dimensional case [3].

The resonance interaction, however, requires the system be prepared in an entangled state and that the coherence of this state be preserved over a sufficiently long time. This quantum correlated state is usually a fragile state, and the coherent superposition can be easily destroyed by the influence of the environment and by spontaneous emission. As far as we know, there are not yet direct observations of this interaction. It is thus a fundamental issue to investigate setups that could allow a measurement of this elusive quantum coherent phenomenon. An appropriate environment around the atoms (or quantum dots), preserving the correlated atomic state for a sufficiently long time, could be exploited in order to observe and measure the resonance interaction. A somehow related effect, the coherent dipolar-induced exchange of excitation (Förster resonance), has been recently ob-

*Electronic address: valentina.notararigo.14@ucl.ac.uk

[†]Electronic address: roberto.passante@unipa.it

[‡]Electronic address: lucia.rizzuto@unipa.it

served for a system of two Rydberg atoms at a separation of about $15 \mu\text{m}$, that is in the short-distance (essentially electrostatic) regime [18]. Observation of the resonance force, in particular in the retarded large-distance regime, would be an important confirmation of an effect due to a quantum coherence property of the system. Also, it could be relevant in different physical processes. For example, the possible important role of resonant interactions in biological systems has been recently argued [19]. An analogous interaction has been shown for two impurities (adatoms) embedded in a semiconductor quantum wire, mediated by the exchange of an electron in the continuum band of the wire [20].

In a recent paper, the interatomic resonance force was shown to be strongly affected by the presence of a structured environment such as a photonic crystal that changes the photon density of states and the dispersion relation [3]. It was shown that in both cases of a one-dimensional photonic crystal and an isotropic three-dimensional crystal, the interaction energy and force can be significantly enhanced up to a 10^3 if the transition frequency of the two identical atoms is close, but external, to the edges of the photonic bandgap. However, in this case the spontaneous emission rate increases too, making even more fragile the entangled state of the two atoms: from one side the resonance force can be much more intense than in free space, but, on the other side, the correlated state decays quicker.

In this Letter, we consider the case of two identical atoms in a generic environment with a photonic bandgap, and with the atomic transition frequency inside the bandgap in the proximity of its edge, assuming a quadratic dispersion relation (effective mass approximation). In the bandgap the photonic density of states is strongly reduced and therefore the spontaneous decay is inhibited (see Refs. [21–23] for the case of a photonic crystal). Inhibited decay rates of spontaneous emission have been recently observed for a quantum dot in a three-dimensional photonic crystal [24]. Several different engineered environments exhibit a photonic bandgap, for example an array of coupled optical cavities, a photonic crystal or a photonic crystal waveguide (for a review, see [25]). The relevance of the configuration here considered is that the spontaneous emission of the atoms is suppressed due to the vanishing photon density of states inside the gap, and this allows to maintain the correlated state for a longer time, hopefully enough to observe the resonance interaction. On the other hand, we will find that the resonance interaction, although decaying faster with the distance compared to the free-space case in the long-distance regime, remains significant and much larger than the dispersion interaction between two atoms in a factorized state (several orders of magnitude). We will show that in the asymptotic limit it decreases as r^{-1} rather than a pure oscillation in the one-dimensional case, while in the three-dimensional case it scales as r^{-2} rather than the r^{-1} free-space behavior. All this indicates that this configuration could be an appropriate

setup for the observation of the resonance interaction. Indications of suppression of dipole-dipole interactions and energy transfer in this case have been discussed in the literature [26], but the results were controversial [27–29]. Also, the possibility of controlling the entanglement of two qubits by coupling to a common photonic bandgap environment has been investigated [30].

The Hamiltonian of our system in the multipolar coupling scheme, within dipole approximation, is

$$\begin{aligned} H &= H_0 + H_I, \\ H_0 &= H_A + H_B + \sum_{\mathbf{k}j} \hbar\omega_k a_{\mathbf{k}j}^\dagger a_{\mathbf{k}j}, \\ H_I &= -i \sum_{\mathbf{k}j} \sqrt{\frac{2\pi\hbar\omega_k}{\Omega}} \mathbf{p}_A \cdot \hat{\mathbf{e}}_{\mathbf{k}j} (a_{\mathbf{k}j} - a_{\mathbf{k}j}^\dagger) \\ &\quad - i \sum_{\mathbf{k}j} \sqrt{\frac{2\pi\hbar\omega_k}{\Omega}} \mathbf{p}_B \cdot \hat{\mathbf{e}}_{\mathbf{k}j} (a_{\mathbf{k}j} e^{i\mathbf{k}\cdot\mathbf{r}} - a_{\mathbf{k}j}^\dagger e^{-i\mathbf{k}\cdot\mathbf{r}}) \end{aligned} \quad (1)$$

where A, B indicate the two atoms with positions $\mathbf{0}$ and $\mathbf{r}$, respectively, the polarization unit vectors $\hat{\mathbf{e}}_{\mathbf{k}j}$ are assumed real, and $\Omega = L(V)$ in the one-dimensional (three-dimensional) case, L and V being respectively the quantization length and the quantization volume. $\mathbf{p}_A$ and $\mathbf{p}_B$ are the atomic operators coupled to the field: in one dimension they have the dimension of a dipole moment per unit length, and in three dimensions they are the atomic dipole moment operators $\boldsymbol{\mu}_A = e\mathbf{r}_A$ and $\boldsymbol{\mu}_B = e\mathbf{r}_B$. In the one-dimensional case the wavevector $\mathbf{k}$ and the position $\mathbf{r}$ are both along the same direction. At the end we will take the continuum limit $\sum_{\mathbf{k}} \rightarrow (L/2\pi) \int dk$ and $\sum_{\mathbf{k}} \rightarrow (V/(2\pi)^3) \int dk k^2 d\Omega_{\mathbf{k}}$ for the one- and three-dimensional cases, respectively. In the dipolar coupling scheme (1), static interatomic dipole-dipole interactions are already included in the interaction with the transverse field and thus mediated by photons [31]. The presence of an external environment is taken into account through the specific photon dispersion relation and density of states.

We assume the two identical atoms prepared in their correlated symmetric or antisymmetric state

$$|\psi\rangle_{\pm} = \frac{1}{\sqrt{2}} (|g_A, e_B; 0_{\mathbf{k}j}\rangle \pm |e_A, g_B; 0_{\mathbf{k}j}\rangle), \quad (2)$$

where g (e) indicates the ground (excited) state of the atom, and $0_{\mathbf{k}j}$ is the photon vacuum. The two states (2) are degenerate and perturbation theory for degenerate states should be used. However, from the expressions of second-order energy corrections for degenerate states [32], it is possible to see that the symmetric and antisymmetric subspaces do not mix each other, provided the dipole matrix elements of the two atoms are equal and the photon dispersion relation is symmetric for $\mathbf{k}$ and $-\mathbf{k}$. If these conditions are verified, the degenerate symmetric and antisymmetric states can be then treated separately and the second-order energy shift is given by nondegenerate perturbation theory. These conditions are

indeed verified in our case, because of our dispersion relation and $\mu_A^{ge} = \mu_B^{ge}$.

The second-order energy shift is then given by

$$\begin{aligned} & \sum_i \frac{\langle \psi^\pm | H_I | i \rangle \langle i | H_I | \psi^\pm \rangle}{E_{\psi^\pm} - E_i} \\ &= \pm \Re \sum_i \frac{\langle g_A, e_B; 0_{\mathbf{k}j} | H_I | i \rangle \langle i | H_I | e_A, g_B; 0_{\mathbf{k}j} \rangle}{\hbar\omega_a - E_i} \\ &+ (L.S.), \end{aligned} \quad (3)$$

where $\Re$ denotes the real part, $|i\rangle$ are intermediate states with energy E_i , $\hbar\omega_a = E_{\psi^\pm}$ is the energy of the atomic excited state, and the $+$ or $-$ sign refers to the symmetric or antisymmetric state in (2), respectively; $(L.S.)$ indicates energy corrections independent from the distance between the two atoms, i.e. the Lamb shifts of the individual atoms. Eq. (3) shows that the resonant interaction energy is strictly related to the resonant energy transfer, that is the excitation transfer between two atoms, whose probability is exactly given by the quantity in the second line of (3), excluding the $\pm$ sign [13]. Therefore our results will be also relevant for the resonant energy transfer.

We are considering the case of an atomic transition frequency inside the photon gap, even if in its proximity, so that the atom-field interaction can be described by second-order perturbation theory (the presence of the photonic gap reduces the resonant pole contribution). There are two possible intermediate states in (3), $|g_A, g_B; 1_{\mathbf{k}j}\rangle$, $|e_A, e_B; 1_{\mathbf{k}j}\rangle$, involving one real or one virtual photon, respectively. We will keep both contributions.

Neglecting distance-independent terms, that do not contribute to the interatomic interaction, we obtain for the second-order energy shift

$$\begin{aligned} \Delta E &= \pm \Re \sum_{\mathbf{k}j} \frac{2\pi\omega_k}{\Omega} (\hat{e}_{\mathbf{k}j})_\ell (\hat{e}_{\mathbf{k}j})_m (p_A^{ge})_\ell (p_B^{eg})_m \\ &\times \left[\frac{e^{i\mathbf{k}\cdot\mathbf{r}}}{ck_a - \omega_k} - \frac{e^{-i\mathbf{k}\cdot\mathbf{r}}}{ck_a + \omega_k} \right], \end{aligned} \quad (4)$$

where $\omega_a = ck_a$ is the atomic transition frequency, $\mathbf{p}_{A,B}^{ge}$ are matrix elements of the atomic dipole operators between ground and excited state.

In the one-dimensional case, after polarization sum and the continuum limit ($L \rightarrow \infty$), we obtain

$$\begin{aligned} \Delta E_{1D} &= \pm 2 |p^{eg}|_\perp^2 \left[\int_0^\infty dk \frac{\omega_k}{ck_a - \omega_k} \cos(kr) \right. \\ &\quad \left. - \int_0^\infty dk \frac{\omega_k}{ck_a + \omega_k} \cos(kr) \right], \end{aligned} \quad (5)$$

where r is the distance between the atoms and subscript $\perp$ indicates the component in the plane orthogonal to the interatomic distance. Similarly, in the three-dimensional

case, after summing over polarizations, taking the continuum limit ($V \rightarrow \infty$) and the angular integration, we obtain

$$\begin{aligned} \Delta E_{3D} &= \pm \frac{1}{\pi} (\mu_A^{ge})_\ell (\mu_B^{eg})_m (-\nabla^2 \delta_{\ell m} + \nabla_\ell \nabla_m) \frac{1}{r} \\ &\times \left[\int_0^\infty dk \frac{\omega_k}{(ck_a - \omega_k)} \frac{\sin(kr)}{k} \right. \\ &\quad \left. - \int_0^\infty dk \frac{\omega_k}{(ck_a + \omega_k)} \frac{\sin(kr)}{k} \right]. \end{aligned} \quad (6)$$

Up to this point, we have not yet specified the environment around the two atoms, except that ω_k depends only on $k = |\mathbf{k}|$. We now assume that the two atoms are embedded in an environment with a photonic band gap with a quadratic dispersion relation between frequency and wavevector, outside the gap and in proximity of its edge. Such a quadratic dispersion relation is typical of a one-dimensional [21, 22] or isotropic three-dimensional [23] photonic crystal in the effective mass approximation, a coupled cavities array [25, 33–35], or an impurity in a quantum wire near the gap [3, 36]. Such a dispersion relation is commonly used to investigate and control radiative properties of atoms or quantum dots in a structured environment [2, 25, 37, 38].

Fig. 1 shows the physical situation we are considering, with the atomic frequency $\omega_a = ck_a$ just above the lower edge ω_v of the gap (for a one-dimensional photonic crystal, the first gap occurs at $k_0 = \pi/\ell$, where ℓ is the periodicity of the crystal [22]) and well below its upper edge ω_c . Near the lower edge of the gap at $k = k_0$, the quadratic dispersion relation is

$$\omega_k = \omega_v - A(k - k_0)^2, \quad (7)$$

where A is a constant that depends on the physical parameters of the structured environment. In the same approximation, the dispersion relation in the region near the upper edge of the gap is $\omega_k = \omega_c + A(k - k_0)^2$.

In the integrals over k in (5) and (6), all values of k should be in principle included. However, only a much smaller range of k gives in our case a significant contribution, allowing a considerable simplification of our calculation. The density of states vanishes inside the gap (that is for photon frequencies between ω_v and ω_c), while it diverges just outside both edges of the gap due to the parabolic dispersion relation (7) [21, 25]. If the atomic frequency ck_a is inside the gap, $\omega_v < ck_a < \omega_c$, and very close to the lower edge ω_v ($ck_a - \omega_v \ll \Delta\omega = \omega_c - \omega_v$), only modes with a frequency just below the gap's lower edge give a significant contribution to the integrals, because of both the large density of states and a near-resonance effect. This can be verified by a direct comparison between the contribution of field modes with $0 \leq k \leq k_0$ and the contribution of modes in a frequency range just above k_0 , for example $k_0 \leq k \leq 3k_0/2$. We now show that the latter contribution is quite smaller if the above conditions are verified. Moreover, this makes




FIG. 1: (color online) The figure shows the position of the atomic frequency $\omega_a = ck_a$, in relation to the lower and upper edge of the first photonic gap at wavenumber k_0 , respectively given by ω_v and ω_c . $\Delta\omega = \omega_c - \omega_v$ is the width of the photonic gap.

clear that the contribution of the remaining modes with $k > 3k_0/2$ will be negligible too.

In the integration range $0 \leq k \leq k_0$ we use the dispersion relation (7); even if this approximation is valid only in the proximity of the gap edge, we can safely extend it to lower values of k because these modes are off-resonance and with a lower density of states, so that they do not significantly contribute to the integration. In the one-dimensional case, we have the following contribution from the k integral in (5)

$$I_{1D} = \int_0^{k_0} dk \left[\frac{\omega_v - A(k - k_0)^2}{ck_a - \omega_v + A(k - k_0)^2} - \frac{\omega_v - A(k - k_0)^2}{ck_a + \omega_v - A(k - k_0)^2} \right] \cos(kr), \quad (8)$$

while, for the three-dimensional case, the k integral in (6) is

$$I_{3D} = \int_0^{k_0} dk \left[\frac{\omega_v - A(k - k_0)^2}{ck_a - \omega_v + A(k - k_0)^2} - \frac{\omega_v - A(k - k_0)^2}{ck_a + \omega_v - A(k - k_0)^2} \right] \frac{\sin(kr)}{k}. \quad (9)$$

We now consider the large-distance limit $r \gg k_0^{-1} \simeq k_a^{-1}$. The parameters k_0, A, ω_v in the integrals above depend on the specific environment and appear in the quadratic dispersion relation (7) valid for $k \sim k_0$ with $k < k_0$; in order to obtain a good interpolation of the dispersion relation for very small values of k where $\omega_k \sim ck$ (because the propagation of a photon with a wavelength much larger than the environment periodicity is that for free space), we can put $\omega_v = Ak_0^2$. In this case, performing the integrals, and taking also into account that in the three-dimensional case space derivatives must be applied to the result (see Eq. (6)), we obtain

$$I_{1D} \simeq \Gamma_1 \frac{\sin(k_0 r)}{r}, \quad (10)$$

where

$$\Gamma_1 = 2 \left(\frac{ck_a Ak_0^2}{c^2 k_a^2 - A^2 k_0^4} + 1 \right) > 0, \quad (11)$$

and

$$I_{3D} \simeq -\Gamma_3 \frac{\cos(k_0 r)}{k_0 r}, \quad (12)$$

where

$$\Gamma_3 = \frac{2ck_a Ak_0^2}{c^2 k_a^2 - A^2 k_0^4} + \sqrt{\frac{Ak_0^2}{ck_a + Ak_0^2}} > 0. \quad (13)$$

We can now compare the contributions obtained above with those coming from the integration above the (first) gap, for example between the first and the second gap in the case of a photonic crystal. For $k_0 < k < 2k_0$, we expect that ranges of k close to k_0 give a quite larger contribution with respect to ranges close to $2k_0$, being closer to resonance. Hence we may evaluate, for both one- and three-dimensional cases, the contribution of k from k_0 to $3k_0/2$, using the dispersion relation in the effective mass approximation above the gap, $\omega_k = \omega_c + A(k - k_0)^2$. The relative k integrals can be evaluated analytically in terms of the system parameters k_a, k_0, A, ω_v . We can meaningfully use typical values for the atomic transition frequency in the optical region and a typical photonic crystal for the environment. Typical numerical values we have used for this check are $\omega_a = ck_a = 2.60 \cdot 10^{15} \text{s}^{-1}$, $\omega_v = Ak_0^2 = 2.58 \cdot 10^{15} \text{s}^{-1}$, $\omega_c = 2.75 \cdot 10^{15} \text{s}^{-1}$; for such values the width of the photonic energy gap $\Delta\omega = \omega_c - \omega_v$ is such that $\omega_a - \omega_v \ll \Delta\omega$, as we have assumed. Using these numerical values, we obtain that the contribution from the range $0 \leftrightarrow k_0$ given by Eqs. (10) and (12) for one and three dimensions respectively, is one order of magnitude larger than the corresponding contribution from the range $k_0 \leftrightarrow 3k_0/2$. This allows us to neglect the latter contribution. Moreover, this gives a strong indication that also all contributions from the ranges $k > 3k_0/2$ will be *a fortiori* negligible, being still more distant from resonance.

Thus, in the one-dimensional case, substitution of (10) and (8) into (5) yields, in the long-distance limit,

$$\Delta E_{1D} \simeq \pm 2\Gamma_1 |p^{eg}|_{\perp}^2 \frac{\sin(k_0 r)}{r}. \quad (14)$$

In the three-dimensional case, in the long-distance limit, substitution of (12) and (9) into (6) yields

$$\begin{aligned} \Delta E_{3D} &\simeq \mp \frac{\Gamma_3}{\pi k_0} (\mu_A^{ge})_{\ell} (\mu_B^{eg})_m (-\nabla^2 \delta_{\ell m} + \nabla_{\ell} \nabla_m) \frac{\cos(k_0 r)}{r^2} \\ &= \mp \frac{\Gamma_3 k_0}{\pi} (\mu_A^{ge})_{\ell} (\mu_B^{eg})_m (\delta_{\ell m} - \hat{r}_{\ell} \hat{r}_m) \frac{\cos(k_0 r)}{r^2}. \end{aligned} \quad (15)$$

In a quasi-static approach, the force between the two atoms can be obtained by taking the opposite of the

derivative of the energy shifts (14) and (15) with respect to the interatomic distance.

The results above should be compared with the analogous quantities obtained for atoms in the free space

$$\Delta E_{1D}^{f.s.} \simeq \pm 2\pi |p^{eg}|_{\perp}^2 k_a \sin(k_a r), \quad (16)$$

[3] and

$$\Delta E_{3D}^{f.s.} \simeq \mp k_a^2 (\mu_A^{ge})_{\ell} (\mu_B^{eg})_m (\delta_{\ell m} - \hat{r}_{\ell} \hat{r}_m) \frac{\cos(k_a r)}{r}. \quad (17)$$

[8, 13].

Equations (14) and (15) are our main results. They show that for $r \gg k_a^{-1} \simeq k_0^{-1}$ the resonant interaction scales with the interatomic distance as r^{-1} in the one-dimensional case and as r^{-2} in the isotropic three-dimensional case. In both cases an oscillation is present, giving a spatially periodic change of the resonance force from attractive to repulsive. Because in free space the respective behaviors are a pure oscillation and r^{-1} (see Eqs. (16) and (17)), our results clearly show that the bandgap environment can qualitatively change the scaling of the interaction with the distance. Although in either cases the resonance force decreases more rapidly with the distance, it must be stressed that the spontaneous decay of the atoms is strongly suppressed by the presence of the environment (see also [39, 40]), preserving the quantum coherence of the correlated state from the decoherence effects due to spontaneous emission.

Although Eqs. (14), (15), (16), (17) show that the environment chosen can reduce the resonance interaction for $k_a r \gg 1$, in a possible experiment to observe the interaction this effect is sensibly overbalanced by the much stronger stability of the correlated state, allowing us to observe the system for a much longer time. This could be an essential point in designing an experiment for a direct observation of the resonance force. Even more, using the typical values of the system parameters given after Eq. (13), our results show that the bandgap environment increases the interaction for distances up to $\sim 40k_a^{-1}$ (it however decreases with respect to the free-space case at larger distances).

To conclude, we have considered the resonance interaction energy between two entangled atoms interacting with the electromagnetic field in the vacuum state, one excited and the other in its ground state, prepared in

their symmetric or antisymmetric state. The two atoms are placed in a structured environment with a photonic band gap and a parabolic dispersion relation, with their transition frequency inside the gap in the proximity of the band edge. We have explicitly considered a one-dimensional system as well an isotropic tridimensional one. We have stressed the formal similarity of the expression obtained for the interatomic interaction energy with the excitation transfer between atoms, allowing a direct extension of our results to this problem too. In our case the spontaneous decay of the atoms is strongly suppressed due to the small photon density of states inside the gap. We have shown that the presence of the environment changes the asymptotic distance dependence of the resonance interaction energy from a pure oscillation to $1/r$ in a one-dimensional system, and from $1/r$ to $1/r^2$ for an isotropic three-dimensional system. This shows explicitly the possibility to control the resonance interaction and force through the environment. On the other hand, even if the resonance force between the two atoms, in the cases considered in this Letter, can be reduced with respect to the free space case for very large distances (but it is still strong with respect to other long-range interactions such as dispersion interactions, even when factorized excited states are considered), the decoherence effect of the spontaneous decay is significantly reduced too. Strong inhibition of the spontaneous emission rate by structured environments such as photonic crystals of about two orders of magnitude has been recently measured [40]. Thus, due to the environment, the quantum coherent atomic superposition can live for a quite long time, of the same order of the lifetime of the excited state in the environment considered, allowing possible detection of this elusive coherent quantum effect. Our results are thus relevant for a future experiment aiming to a direct measurement of the resonance interaction and force, using atoms or quantum dots embedded in an appropriate structured environment with a photonic bandgap.

Acknowledgments

The authors gratefully acknowledge financial support by the Julian Schwinger Foundation and by MIUR.

-
- [1] A. Salam, *Molecular quantum electrodynamics in the Heisenberg picture: A field theoretic viewpoint*, Int. Rev. Phys. Chem. **27**, 405 (2008).
 - [2] R. El-Ganainy and S. John, *Resonant dipole-dipole interaction in confined and strong-coupling geometries*, New J. Phys. **15**, 083033 (2013).
 - [3] R. Incardone, T. Fukuta, S. Tanaka, T. Petrosky, L. Rizzuto, and R. Passante *Enhanced resonant interaction between two entangled identical atoms in a photonic crystal*,

Phys. Rev. A **89**, 062117 (2014).

- [4] D.L. Andrews and J.S. Ford, *Resonant energy transfer: Influence of neighboring matter absorbing in the wavelength of the acceptor*, J. Chem. Phys. **139**, 014107 (2013).
- [5] G.D. Scholes, *Long-range resonance energy transfer in molecular systems*, Annu. Rev. Phys. Chem. **54**, 57 (2003).
- [6] A. Olaya-Castro, A.N. Nazir, and G.R. Fleming,

- Quantum-coherent energy transfer: implications for biology and new energy technologies*, Phil. Trans. R. Soc. A **370**, 3613 (2012).
- [7] G. Compagno, R. Passante, and F. Persico, *Atom-field interactions and dressed atoms*, Cambridge University Press, Cambridge, UK 1995.
 - [8] D.P. Craig and T. Thirunamachandran, *Molecular Quantum Electrodynamics*, Dover Publ., Mineola, New York, 1998.
 - [9] M. Donaire, R. Guérout, A. Lambrecht, *Quasiresonant van der Waals interaction between nonidentical atoms*, Phys. Rev. Lett. **115**, 033201 (2015).
 - [10] P. R. Berman, *Interaction energy of nonidentical atoms*, Phys. Rev. A **91**, 042127 (2015).
 - [11] P. Barcellona, R. Passante, L. Rizzuto, and S.Y. Buhmann, *van der Waals interactions between excited atoms in generic environments*, Phys. Rev. A, in press (2016).
 - [12] P.W. Milonni and S.M.H. Rafsanjani, *Distance dependence of two-atom dipole interactions with one atom in an excited state*, Phys. Rev. A **92**, 062711 (2015).
 - [13] A. Salam *Molecular Quantum Electrodynamics*, John Wiley & Sons, Hoboken, New Jersey, 2010.
 - [14] R. Messina, R. Passante, L. Rizzuto, S. Spagnolo, and R. Vasile, *Casimir-Polder forces, boundary conditions and fluctuations*, J. Phys. A **41**, 164031 (2008).
 - [15] F. Intravaia, C. Henkel, and M. Antezza, in: *Casimir Physics*, edited by D. Dalvit, P. Milonni, D. Roberts, and F. Rosa, Lecture Notes in Physics 834, Springer, Berlin 2011, p. 345.
 - [16] M.A. Wilson, P. Bushev, J. Eschner, F. Schmidt-Kaler, C. Becher, R. Blatt, and U. Dorner, *Vacuum-field level shifts in a single trapped ion mediated by a single distant mirror*, Phys. Rev. Lett. **91**, 213602 (2003).
 - [17] R. Passante, F. Persico, and L. Rizzuto *Spatial correlations of vacuum fluctuations and the Casimir-Polder potential*, Phys. Lett. A **316**, 29 (2003).
 - [18] S. Ravets, H. Labuhn, D. Barredo, L. Béguin, T. Lahaye, and A. Browaeys, *Coherent dipole-dipole coupling between two single Rydberg atoms at an electrically-tuned Förster resonance*, Nature Phys. **10**, 914 (2014).
 - [19] J. Preto, M. Pettini, and J.A. Tuszynski, *Possible role of electrodynamical interactions in long-distance biomolecular recognition*, Phys. Rev. E **91**, 052710 (2015).
 - [20] S. Tanaka, R. Passante, T. Fukuta, and T. Petrosky, *Nonperturbative approach for the electronic Casimir-Polder effect in a one-dimensional semiconductor*, Phys. Rev. A **88**, 022518 (2013).
 - [21] S. John and J. Wang, *Quantum electrodynamics near a photonic band gap: photon bound states and dressed atoms*, Phys. Rev. Lett. **64**, 2418 (1990).
 - [22] D.G. Angelakis, P.L. Knight, and E. Paspalakis, *Photonic crystals and inhibition of spontaneous emission: an introduction*, Contemp. Phys. **45**, 303 (2004).
 - [23] S. John and J. Wang, *Quantum optics of localized light in a photonic band gap*, Phys. Rev. B **43**, 12772 (1991).
 - [24] M.D. Leistikow, A.P. Mosk, E. Yeganegi, S.R. Huisman, A. Lagendijk, and W.L. Vos, *Inhibited Spontaneous Emission of Quantum Dots Observed in a 3D Photonic Band Gap*, Phys. Rev. Lett. **107**, 193903 (2011).
 - [25] P. Lodahl, S. Mahmoodian, and S. Stobbe, *Interfacing single photons and single quantum dots with photonic nanostructures*, Rev. Mod. Phys. **87**, 347 (2015).
 - [26] G. Kurizki and A.Z. Genack, *Suppression of molecular interactions in periodic dielectric structures*, Phys. Rev. Lett. **61**, 2269 (1988).
 - [27] Q. Zheng, T. Kobayashi, and T. Sekiguchi, *Comment on “Suppression of Molecular Interactions in Periodic Dielectric Structures”*, Phys. Rev. Lett. **77**, 406 (1996).
 - [28] G. Kurizki, A.G. Kofman, and A.Z. Genack, *Reply to the Comment by Qianbing Zheng, Takayoshi Kobayashi, and Takashi Sekiguchi*, Phys. Rev. Lett. **77**, 407 (1996).
 - [29] G. Kurizki, *Two-atom resonant radiative coupling in photonic band structures*, Phys. Rev. A **42**, 2915 (1990).
 - [30] J. Wang, L. Jiang, H. Zhang, T.H. Huang, H.Z. Zhang, *Controlled entanglement of two atoms in photonic crystals*, Opt. Comm. **284**, 5323 (2011).
 - [31] E.A. Power and S. Zienau, *Coulomb gauge in non-relativistic quantum electrodynamics and the shape of spectral lines*, Phil. Trans. Royal Soc. A **251**, 54 (1959).
 - [32] L.I. Schiff, *Quantum Mechanics*, McGraw Hill Book Company, Singapore 1968.
 - [33] A. Goban, C.L. Hung, S.P. Yu, J.D. Hood, J.A. Muniz, J.H. Lee, M.J. Martin, A.C. McClung, K.S. Choi, D.E. Chang, O. Painter, and H.J. Kimble, *Atom-light interactions in photonic crystals*, Nat. Commun. **5**, 3808 (2014).
 - [34] M.J. Hartmann, F.G.S.L. Brandão, and M.B. Plenio, *Quantum many-body phenomena in coupled cavity arrays*, Laser Photonics Rev. **2**, 527 (2008).
 - [35] J.S. Douglas, H. Habibian, C.L. Hung, A.V. Gorshkov, H.J. Kimble, and D.E. Chang, *Quantum many-body models with cold atoms coupled to photonic crystals*, Nature Photonics **9**, 326 (2015).
 - [36] S. Tanaka, S. Garmon, and T. Petrosky, *Nonanalytic enhancement of the charge transfer from adatom to one-dimensional semiconductor superlattice and optical absorption spectrum*, Phys. Rev. B **73**, 115340 (2006).
 - [37] S. John and T. Quang, *Spontaneous emission near the edge of a photonic band gap*, Phys. Rev. A **50**, 1764 (1994).
 - [38] X.H. Wang, B.Y. Gu, R. Wang, and H.Q. Xu, *Decay kinetic properties of atoms in photonic crystals with absolute gaps*, Phys. Rev. Lett. **91**, 113904 (2003).
 - [39] P. Lodahl, A.F. van Driel, I.S. Nikolaev, A. Irman, K. Overgaag, D. Vanmaekelbergh, and W.L. Vos, *Controlling the dynamics of spontaneous emission from quantum dots by photonic crystals*, Nature **430**, 654 (2004).
 - [40] Q. Wang, S. Stobbe, P. Lodahl, *Mapping the local density of optical states of a photonic crystal with single quantum dots*, Phys. Rev. Lett. **107**, 167404 (2011).